

# HIPOTERMIA PERIOPERATORIA

José Ignacio Redondo García

DVM, PhD, MSc (Statistics)  
Universidad CEU Cardenal Herrera



**B|BRAUN**  
SHARING EXPERTISE



**CEU** | *Universidad  
Cardenal Herrera*



# Córdoba logra un nuevo récord de temperatura máxima en España desde que hay registros, con 46,9°C

OLA DE CALOR >

## España bate su récord de temperatura: 46,9 grados

EL TIEMPO ANÁLISIS


Córdoba supera su temperatura máxima histórica



AEMET   
@AEMET\_Esp

Seguir

13/07 23:31 **#AEMET** **#avisos** nivel rojo por temp. max para mañana en Andalucía +info [goo.gl/U8fS](https://goo.gl/U8fS)



23:31 - 13 jul. 2017

4 Retweets 10 Me gusta





¿POR QUÉ?



## ***Complications and Treatment of Mild Hypothermia***

*Daniel I. Sessler, M.D.\**

Even mild hypothermia produces marked postoperative thermal discomfort.<sup>1,73</sup> Patients often indicate that feeling cold in the immediate postoperative period is the worst part of their hospitalization, sometimes rating it worse than surgical pain. Given the appropriate efforts to treat surgical pain, it would similarly seem appropriate to prevent and treat thermal discomfort.<sup>20</sup>



# monitorización






# monitorización

## Comparison of three methods of temperature measurement in hypothermic, euthermic, and hyperthermic dogs

Rebecca J. Greer, DVM; Leah A. Cohn, DVM, PhD, DACVIM; John R. Dodam, DVM, PhD, DACVA;  
Colette C. Wagner-Mann, DVM, PhD; F. A. Mann, DVM, MS, DACVS, DAVECC

**Results**—Variability between duplicate near-simultaneous temperature measurements was greatest for the auricular thermometer and least for the TTPA catheter. Measurements obtained by use of the rectal thermometer were in closest agreement with core body temperature; for all other devices, temperature readings typically underestimated core body temperature.

**Conclusions and Clinical Relevance**—Among the 3 methods of temperature measurement, rectal thermometry provided the most accurate estimation of core body temperature in dogs. (*J Am Vet Med Assoc* 2007;230:1841–1848)





# Perioperative hypothermia

Stacy R. Armstrong, DVM, Brian K. Roberts, DVM, DACVECC and Michael Aronsohn, VMD, DACVS

Hypothermia is defined as a decrease in normal body temperature below 37 °C (99 °F).<sup>1</sup> The normal canine and feline temperature range is 37.8–39.2 °C (100.0–102.5 °F). Mild hypothermia in veterinary patients has been defined as body temperature between 32 and 37 °C (90 and 99 °F), while moderate hypothermia is temperatures between 28 and 32 °C (82 and 90 °F). Severe hypothermia is defined as temperatures below 28 °C (82 °F).<sup>2,3</sup>

| HIPOTERMIA          | TEMPERATURA   |
|---------------------|---------------|
| Hipertermia         | > 39,50°C     |
| Normotermia         | 38,50-39,50°C |
| Hipotermia ligera   | 38,49-36,50°C |
| Hipotermia moderada | 36,49-34,00°C |
| Hipotermia grave    | <34,00°C      |





10.1136/vr.100476 | **Veterinary Record**

# Retrospective study of the prevalence of postanaesthetic hypothermia in dogs

J.I. Redondo, P. Suesta, I. Serra, C. Soler, G. Soler, L. Gil, R. J. Gómez-Villamandos

10.1136/vr.100184 | **Veterinary Record**

# Retrospective study of the prevalence of postanaesthetic hypothermia in cats

J. I. Redondo, P. Suesta, L. Gil, G. Soler, I. Serra, C. Soler



# perros


## incidencia

hipotermia leve 51,5 %

hipotermia moderada 29,3 %

hipotermia grave 2,7 %

83,5%






# gatos

## incidencia


hipotermia leve 26,5 %  
hipotermia moderada 60,4 %  
hipotermia grave 10,5 %

97,4%





# termorregulación





# mecanismos intercambio térmico

convección

radiación

conducción

evaporación




FIGURA 1. Formas de intercambio térmico con el medio.



# causas

Anesthesiology  
2000; 92:578-96  
© 2000 American Society of Anesthesiologists, Inc.  
Lippincott Williams & Wilkins, Inc.

## ***Perioperative Heat Balance***

*Daniel I. Sessler, M.D.\**


- ▶ acción de los anestésicos sobre SNC
- ▶ vasodilatación periférica
- ▶ condiciones ambientales






## causas

### ► manejo quirúrgico



- desinfectantes
- apertura cavidades
- tiempo quirúrgico





# consecuencias

- ▶ **METABÓLICAS**
  - ▶ disminución en la metabolización de los anestésicos
  - ▶ hiperglucemia inicial ⇒ hipoglucemia
- ▶ **ELECTROLÍTICAS**
  - ▶ hipokalemia
  - ▶ hipernatremia
  - ▶ hipocalcemia
- ▶ **HIPOPROTEINEMIA**
- ▶ **CARDIOVASCULARES**
  - ▶ retraso tiempo de coagulación
  - ▶ aumento viscosidad sanguínea
  - ▶ formación de coágulos y CID
  - ▶ fibrilación ventricular
- ▶ **RESPIRATORIAS**
  - ▶ bradipnea
  - ▶ hipoxia tisular
  - ▶ metabolismo anaerobio
  - ▶ acidosis láctica
- ▶ **INMUNOLÓGICAS**
  - ▶ hipoxia tisular ▶ muerte neutrófilos y granulocitopenia
  - ▶ retraso en la cicatrización
  - ▶ aumento incidencia infecciones postquirúrgicas



# fases

**1** fase

1 hora de duración.  
Temperatura corporal  
disminuye de 1°C a  
1,5°C

**2** fase

2-3 horas de duración.  
Pérdida de temperatura  
lenta y de forma lineal


**3** fase

3ª o 4ª hora de anestesia.  
La temperatura corporal  
permanece más o menos  
constante.

Anesthesiology  
2000; 92:578-96  
© 2000 American Society of Anesthesiologists, Inc.  
Lippincott Williams & Wilkins, Inc.

## *Perioperative Heat Balance*

Daniel I. Sessler, M.D.\*






# perros

## ► evolución de la temperatura

► T<sup>a</sup> basal media →  $38,7 \pm 0,6$  °C

► T<sup>a</sup> intraoperatoria media →  $36,8 \pm 1,4$  °C






# perros

## ► evolución de la temperatura

► T<sup>a</sup> basal media →  $38,7 \pm 0,6$  °C

► T<sup>a</sup> intraoperatoria media →  $36,8 \pm 1,4$  °C






# perros

## ► evolución de la temperatura

► T<sup>a</sup> basal media →  $38,7 \pm 0,6$  °C

► T<sup>a</sup> intraoperatoria media →  $36,8 \pm 1,4$  °C






# perros

## ► evolución de la temperatura

► T<sup>a</sup> basal media →  $38,7 \pm 0,6$  °C

► T<sup>a</sup> intraoperatoria media →  $36,8 \pm 1,4$  °C






# gatos

## ► evolución de la temperatura

► T<sup>a</sup> basal media →  $38,2 \pm 1,0$  °C

► T<sup>a</sup> intraoperatoria media →  $35,6 \pm 1,5$  °C






# gatos

## ► evolución de la temperatura

► T<sup>a</sup> basal media →  $38,2 \pm 1,0$  °C

► T<sup>a</sup> intraoperatoria media →  $35,6 \pm 1,5$  °C






# gatos

## ► evolución de la temperatura

► T<sup>a</sup> basal media →  $38,2 \pm 1,0$  °C

► T<sup>a</sup> intraoperatoria media →  $35,6 \pm 1,5$  °C






# gatos

## ► evolución de la temperatura

► T<sup>a</sup> basal media →  $38,2 \pm 1,0$  °C

► T<sup>a</sup> intraoperatoria media →  $35,6 \pm 1,5$  °C









# tratamiento

## calentamiento pasivo

- ▶ temperatura ambiente

### THE OPTIMUM TEMPERATURE OF OPERATING-ROOMS

By EMMET RIXFORD, San Francisco

CALIFORNIA AND WESTERN MEDICINE December, 1924

*Brit. J. industr. Med.*, 1963, 20, 284.

### THE CONTROL OF OPERATING-SUITE TEMPERATURES\*

BY


F. P. ELLIS

*British Navy Staff, Washington, D.C.*

### Ambient operating room temperature: mother, baby or surgeon?

P Sultan ✉, A S Habib, B Carvalho

*BJA: British Journal of Anaesthesia*, Volume 119, Issue 4, 1 October 2017, Pages 839,





# tratamiento

## calentamiento pasivo

- Uso de mantas y empapadores





# tratamiento

## calentamiento activo superficial

- ▶ mantas eléctricas
- ▶ bolsas y guantes con agua caliente



JAN

JOURNAL OF ADVANCED NURSING

### REVIEW PAPER

### Effectiveness of cutaneous warming systems on temperature control: meta-analysis

Cristina Maria Galvão, Yuanyuan Liang & [Alexander M. Clark](#)






# tratamiento

## calentamiento activo superficial

- ▶ lámparas infrarrojas
- ▶ aire caliente forzado





# tratamiento

## calentamiento activo central

**Warming of intravenous and irrigation fluids for preventing inadvertent perioperative hypothermia (Review)**

Campbell G, Alderson P, Smith AF, Warttig S

## Heated intravenous fluids alone fail to prevent hypothermia in cats under general anaesthesia

Geraldine Jourdan, Caroline Didier, Erwan Chotard, Sandra Jacques and Patrick Verwaerde

*Journal of Feline Medicine and Surgery*  
1–5  
© The Author(s) 2017  
Reprints and permissions:  
sagepub.co.uk/journalsPermissions.nav  
DOI: 10.1177/1098612X16688990  
journals.sagepub.com/home/jfms

This paper was handled and processed  
by the European Editorial Office (ISFM)  
for publication in *JFMS*



- ▶ lavados abdominales y pleurales.
- ▶ enemas con agua caliente
- ▶ fluido salino caliente en la vejiga

## Effect of peritoneal lavage solution temperature on body temperature in anaesthetised cats and small dogs

D. C. Barnes, E. A. Leece, T. A Trimble, J. L. Demetriou

Veterinary Record (2017)

doi: 10.1136/vr.103894



## „Pre-warming“ – Einführung in ein Verfahren zur Vermeidung der perioperativen Hypothermie

*“Pre-warming” – How can Perioperative Hypothermia be Avoided?*

### CALENTAMIENTO PREVENTIVO

Calentamiento superficial previo  
a la inducción anestésica  
(30 minutos antes)





# Comparison of two methods for the management of intraoperative hypothermia in dogs

J. O. Kibanda, M. Gurney

Veterinary Record (2012) 170, 392a      doi: 10.1136/vr.100690

| TABLE 1: Comparative parameters between the two groups |                         |                   |         |
|--------------------------------------------------------|-------------------------|-------------------|---------|
|                                                        | Forced warm-air blanket | Heated mat        | P Value |
| Temperature at preanaesthetic examination              | 38.6±0.49°C             | 38.3±0.51°C       | 0.0004  |
| First temperature at instrumentation                   | 37.5±0.68°C             | 37.6±0.59°C       | 0.42    |
| Time from induction to draping                         | 79.2±19.8 minutes       | 75.5±20.8 minutes | 0.32    |
| Temperature at draping                                 | 35.9±0.79°C             | 36.3±0.78°C       | 0.005   |
| Rate of heat loss induction to draping                 | 0.02±0.01°C/min         | 0.017±0.01°C/min  | 0.06    |
| Temperature one hour after draping                     | 36.2±0.7°C              | 36.6±0.8°C        | 0.002   |
| Rate of heat gain during first hour of surgery         | 0.004±0.009°C/min       | 0.005±0.007°C/min | 0.6     |
| Temperature nadir                                      | 35.7±0.7°C              | 36.2±0.7°C        | 0.0005  |
| Temperature end of surgery                             | 36.4±0.7°C              | 36.9±0.8°C        | 0.0004  |
| Surgery time                                           | 81±23 minutes           | 88±25 minutes     | 0.1     |



# factores de riesgo

▶perros

## Retrospective study of the prevalence of postanaesthetic hypothermia in dogs

J.I. Redondo, P. Suesta, I. Serra, C. Soler, G. Soler, L. Gil, R. J. Gómez-Villamandos

| TABLE 3: Predictors of temperature (°C) at the end of the procedure in 1525 dogs by multiple regression analysis |                            |                           |                               |         |             |             |
|------------------------------------------------------------------------------------------------------------------|----------------------------|---------------------------|-------------------------------|---------|-------------|-------------|
|                                                                                                                  | Unstandarised coefficients | Standardised coefficients | 95% confidence interval for B |         |             |             |
|                                                                                                                  | B                          | se                        | Beta                          | P value | Lower bound | Upper bound |
| (Constant)                                                                                                       | 20.460                     | 2.373                     |                               | 0       | 15.810      | 25.119      |
| → Baseline temperature                                                                                           | 0.441                      | 0.061                     | 0.152                         | 0       | 0.320       | 0.561       |
| → Body surface                                                                                                   | 1.451                      | 0.112                     | 0.274                         | 0       | 1.232       | 1.670       |
| → Preanaesthetic time                                                                                            | −0.010                     | 0.002                     | −0.120                        | 0       | −0.014      | −0.006      |
| → Duration of anesthesia                                                                                         | −0.010                     | 0.001                     | −0.385                        | 0       | −0.011      | −0.009      |
| → ASA                                                                                                            | −0.122                     | 0.035                     | −0.075                        | 0.001   | −0.191      | −0.052      |
| → Thoracic surgery                                                                                               | −1.645                     | 0.252                     | −0.139                        | 0       | −2.138      | −1.151      |
| → Diagnostic procedures                                                                                          | −0.227                     | 0.098                     | −0.050                        | 0.021   | −0.420      | −0.034      |
| → Dorsal recumbency                                                                                              | −0.507                     | 0.071                     | −0.158                        | 0       | −0.646      | −0.369      |
| → Sternal recumbency                                                                                             | −0.262                     | 0.110                     | −0.052                        | 0.017   | −0.477      | −0.046      |



# recuperación y mortalidad

## ►perros

Relación hipotermia, recuperación anestésica y mortalidad postoperatoria

### Papers

10.1136/vr.100476 | Veterinary Record | 1 of 5

### Retrospective study of the prevalence of postanaesthetic hypothermia in dogs

J.I. Redondo, P. Suesta, I. Serra, C. Soler, G. Soler, L. Gil, R. J. Gómez-Villamandos

TABLE 4: Deaths, mortality index and time to extubation in minutes (mean±sd) by ranks of hypothermia

|                      | N    | Deaths | Mortality index (%) | Time to extubation (minutes) |
|----------------------|------|--------|---------------------|------------------------------|
| Hyperthermia         | 31   | 0      | 0                   | 10±9                         |
| Normothermia         | 220  | 1      | 0.45                | 8±5                          |
| Slight hypothermia   | 785  | 5      | 0.64                | 8±6                          |
| Moderate hypothermia | 447  | 6      | 1.34                | 7±6                          |
| Severe hypothermia   | 42   | 0      | 0                   | 14±21                        |
| Total                | 1525 | 12     | 0.79                | 8±7                          |

N Total number of cases, Deaths Number of dogs that died



# factores de riesgo

## ▶gatos

Papers 10.1136/vr.100184 | Veterinary Record | 1 of 4

Retrospective study of the prevalence of postanaesthetic hypothermia in cats

J. I. Redondo, P. Suesta, L. Gil, G. Soler, I. Serra, C. Soler

TABLE 2: Predictors of temperature (°C) at the end of the procedure in 275 cats by multiple regression analysis

| Variable                            | $\beta$ | P    | 95% CI for $\beta$ |       |
|-------------------------------------|---------|------|--------------------|-------|
|                                     |         |      | Lower              | Upper |
| → Basal temperature                 | 0.14    | 0.02 | 0.05               | 0.53  |
| → Duration of anaesthesia (minutes) | -0.10   | 0.01 | -0.01              | 0.00  |
| → Reason                            |         |      |                    |       |
| ORTH                                | -0.45   | 0.00 | -1.95              | -0.97 |
| ABD                                 | -0.38   | 0.00 | -1.53              | -0.70 |
| → Anaesthetic risk                  |         |      |                    |       |
| High risk                           | -0.14   | 0.02 | -1.10              | -0.24 |

ABD Abdominal surgery, ORTH Orthopaedic surgery,  $\beta$  Standardised coefficients for the studied variables in the multiple regression model



# recuperación y mortalidad

►gatos

Relación hipotermia, recuperación anestésica y mortalidad postoperatoria

Papers 10.1136/vr.100184 | Veterinary Record | 1 of 4

Retrospective study of the prevalence of postanaesthetic hypothermia in cats


J. I. Redondo, P. Suesta, L. Gil, G. Soler, I. Serra, C. Soler

| TABLE 3: Time to extubation in minutes (mean (sd)), deaths and mortality index by ranks of hypothermia in 275 cats undergoing anaesthesia |                              |                |        |                     |
|-------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|----------------|--------|---------------------|
|                                                                                                                                           | Time to extubation (minutes) | Number of cats | Deaths | Mortality index (%) |
| Hyperthermia                                                                                                                              | 23 (0)                       | 2              | 0      | 0                   |
| Normothermia                                                                                                                              | 13 (11)                      | 5              | 0      | 0                   |
| Slight hypothermia                                                                                                                        | 12 (7)                       | 73             | 2      | 2.73                |
| Moderate hypothermia                                                                                                                      | 13 (10)                      | 166            | 2      | 1.20                |
| Severe hypothermia                                                                                                                        | 26 (25)                      | 29             | 2      | 6.89                |
| Total                                                                                                                                     | 14 (12)                      | 275            | 6      | 2.18                |



## conclusiones

- ▶ La hipotermia intraoperatoria y postoperatoria es muy frecuente en **perros (83,6%)** y **gatos (97,4%)**.
- ▶ En el perro los **factores de riesgo** son: el tiempo de inducción, tiempo de anestesia, superficie corporal, sexo, edad, tipo de cirugía, riesgo anestésico y el decúbito adoptado durante la anestesia.
- ▶ En el **gato**, por su parte, son: tiempo de anestesia, sexo, superficie corporal, tipo de cirugía, el riesgo anestésico y el tipo de decúbito.
- ▶ El **tiempo de extubación** aumenta en ambas especies en pacientes con hipotermias graves ( $< 34^{\circ}\text{C}$ ).
- ▶ El índice de **mortalidad** aumenta en gatos con hipotermia grave.






# Córdoba





## Willis Carrier





# HIPOTERMIA PERIOPERATORIA

José Ignacio Redondo García

DVM, PhD, MSc (Statistics)  
Universidad CEU Cardenal Herrera



**B|BRAUN**  
SHARING EXPERTISE



**CEU**

*Universidad  
Cardenal Herrera*